
Socioecological Determinants of Recovery in Opioid Substitution Treatment Patients: A Systematic Review and Thematic Synthesis

Vicky Carlisle, PhD Student, School of Psychological Science & Population Health Sciences





National Drug & Alcohol Treatment Population



Reporting Period


Source: NDTMS – National Drug Treatment Monitoring System (2019)





Treatment completions

Figure 6.1.1 Proportion of exits that are treatment completed free of dependence by the four substance groups 2017-18



Source: NDTMS – National Drug Treatment Monitoring System (2019)





Aims

- Aims:
- Identify facilitators and barriers to recovery within OST
- Familiarisation with qualitative literature on OST
- Identify possible targets for intervention development
- Inclusion criteria:
- Any country (English language papers only)
- Adults (18+)
- Service users and stakeholder views
- Peer reviewed articles published in last 20 years





Characteristics of Included Studies (n=32)

- o UK (10)
- o USA (8)
- o Ireland (5)
- o Canada (3)
- o Australia (2)
- Norway (2)
- o Italy (1)
- Sweden (1)



Service users (24 studies): n=680

Stakeholders (3 studies): n=63

Both (5 studies): n=263

Total participants: n=1,006






Socioecological Model (Bronfenbenner, 1979)













Trauma

[describing experience of children's home] "Life there was horrible, horrible, horrible, some terrible things, Oh I don't even want to talk about them..." (Moran et al, 2018).

"You know, I'm busy. My mother's in the hospital, you know. I'm dealing with the death of a brother and my son was just murdered. I got a lot going on in my life right now." (Lindsey-Smith & Rosen, 2009).

"I lost two children in 9 days. My son, he hung himself in prison, three days left in his sentence. My daughter committed suicide 9 days later. She went out, bought Xanax (alprazolam), and took the whole lot and she took about nine bags of heroin as well and just threw herself off the bridge. Afterwards, I tried to kill myself." (Moran et al, 2018).





Trauma:

ACEs

Bereavement

Violence

Rape

Street sex work





Trauma:

ACEs

Bereavement

Violence

Rape

Street sex work

Drug use (self-medication)








ACEs

Bereavement

Violence

Rape

Street sex work










ACEs

Bereavement

Violence


Rape

Street sex work


















Integrative Epidemiology

University of BRISTOL













Summary

- Recovery from opioid dependency is a complex problem that requires a complex solution.
- Interventions targeted at the individual only will have limited success.
- Possible targets for intervention are trauma, stigma and isolation.











Acknowledgements

Thanks to:

Tobacco and Alcohol Research Group:

Angela Attwood Claire Braboszcz Anna Blackwell Laura Brocklebank Sarah Burrows-Weeks

Vicky Carlisle
Emily Crowe
Katie De-Loyde
Katie Drax
Maddy Dyer
Kayleigh Easey
Andy Eastwood
Jennifer Ferrar
Will Gawned

Gemma Hammerton Abigail Jackson

Elis Haan

Senior Lecturer and TARG co-director

Senior Research Associate Senior Research Associate Research Associate Research Administrator PhD Student PhD Student Research Associate PhD Student PhD Student PhD Student PhD Student

PhD Student Research Associate Administrator PhD Student Research Fellow

Senior Research Associate

Andy Keavey

Rebecca Lawn
Joe Matthews
Liam Mahedy
Osama Mahmoud
Olivia Maynard

Maren Müller-Glodde Marcus Munafò

Hannah Sallis Laura Schellhas Chris Stone Andy Skinner Steph Suddell Jackie Thompson Robert Thibault Sophie Turnbull Robyn Wootton University of BRISTOL



Developer, Cambridge Cognition

PhD Student

Research Associate

Senior Research Associate Senior Research Associate Lecturer and TARG co-director

PhD Student

Professor and TARG co-director Senior Research Associate

PhD Student

Senior Research Associate

Research Fellow PhD Student

Research Associate Research Fellow Research Associate

Senior Research Associate



Contact: Vicky.carlisle@bristol.ac.uk;

@Vic_Carlisle; @BristolTARG; http://www.bristol.ac.uk/targ







References

- Moran, L., Keenan, E., & Elmusharaf, K. (2018). Barriers to progressing through a methadone maintenance treatment programme: perspectives of the clients in the Mid-West of Ireland's drug and alcohol services. BMC Health Serv Res, 18(1), 911.
- PHE. (2018.). Adult substance misuse statistics from the National Drug Treatment Monitoring System (NDTMS). Retrieved from https://assets.publishing.service.gov.uk/government/uploads/system/uplods/attachment_data/file/658056/Adult-statistics-from-the-national-drug treatment-monitoring-system-2016-2017.pdf
- Smith, M. L., & Rosen, D. (2009). Mistrust and self-isolation: Barriers to social support for older adult methadone clients. [References]. Journal of Gerontological Social Work. Vol., 52(7), 653–667.





Community

Caught between two cultures:

"Recovering heroin users want to be normal and yet they also want to retain their individuality, they aspire to the mundane and yet they are anxious that this is also boring, they want to comply with social norms and conventions and yet they want to retain their distinctiveness" (Nettleton, Neale, & Pickering, 2013)







Interpersonal




Influences of family and friends (positive and negative):

"It happened that I started abusing again. When I went back to Terni I found all my friends, the old ones, and I started again right away. Would you believe it! When you fall back into that crowd, they offer it to you right away and it's over." (Cordella, Greco, Elia, & Varazi, 2017).













We have a retention problem too













Intrapersonal

Self-Efficacy & Locus of Control



"Perhaps more significantly, participants framed their recovery in avoidance language (staying away from drugs, walking away from temptation, hiding from trouble) as opposed to engagement language (e.g., finding a job, maintaining a healthy lifestyle)" (Redden, Tracy, & Shafer, 2013)



